

cows; and for advancement of first pubertal estrus in replacement beef heifers.

(B) For synchronization of estrus in lactating dairy cows.

(C) For synchronization of the return to estrus in lactating dairy cows inseminated at the immediately preceding estrus.

(iii) *Limitations.* Do not use in beef or dairy heifers of insufficient size or age for breeding or in animals with abnormal, immature, or infected genital tracts. Do not use in beef cows that are fewer than 20 days postpartum. Do not use an insert more than once. To prevent the potential transmission of venereal and bloodborne diseases, the inserts should be disposed after a single use. Administration of vaginal inserts for periods greater than 7 days may result in reduced fertility. Dinoprost solution provided by No. 000009 in § 510.600(c) of this chapter.

(2) *Ewes*—(i) *Amount.* Administer one intravaginal insert per animal for 5 days.

(ii) *Indications for use.* For induction of estrus in ewes (sheep) during seasonal anestrus.

(iii) *Limitations.* Do not use in animals with abnormal, immature, or infected genital tracts; or in ewes that have never lambed. Do not use an insert more than once. To prevent the potential transmission of venereal and bloodborne diseases, the inserts should be disposed after a single use.

[74 FR 59074, Nov. 17, 2009, as amended at 75 FR 63085, Oct. 14, 2010]

§ 529.2150 Sevoflurane.

(a) *Specifications.* Sevoflurane liquid.

(b) *Sponsors.* See Nos. 000074, 012164, and 066794 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* For induction of surgical anesthesia: up to 7 percent sevoflurane. For maintenance of surgical anesthesia: 3.7 to 4 percent sevoflurane with oxygen in the absence of premedication and 3.3 to 3.6 percent in the presence of premedication.

(2) *Indications for use.* For induction and maintenance of general anesthesia in dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[64 FR 71640, Dec. 22, 1999, as amended at 73 FR 25508, May 7, 2008; 74 FR 10484, Mar. 11, 2009, 75 FR 1021, Jan. 8, 2010; 76 FR 16533, Mar. 24, 2011]

§ 529.2464 Ticarcillin powder.

(a) *Specifications.* Each vial contains ticarcillin disodium equivalent to 6 grams of ticarcillin to be reconstituted with 25 milliliters of sterile water for injection or sterile physiological saline.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 6 grams per day, intrauterine, for 3 consecutive days during estrus.

(2) *Indications for use.* *Horses.* Intrauterine treatment of endometritis caused by beta-hemolytic streptococci.

(3) *Limitations.* For intrauterine use in horses only. Infuse aseptically. Not for use in horses raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37336, Aug. 18, 1992, as amended at 60 FR 55660, Nov. 2, 1995]

§ 529.2503 Tricaine methanesulfonate.

(a) *Chemical name.* Ethyl-*m*-aminobenzoate methanesulfonate.

(b) *Sponsor.* See Nos. 050378 and 051212 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used for the temporary immobilization of fish, amphibians, and other aquatic cold-blooded animals (poikilotherms) as an aid in handling during manual spawning (fish stripping), weighing, measuring, marking, surgical operations, transport, photography, and research.

(2) It is used as follows:

(i) For fish the drug is added to ambient water at a concentration of from 15 to 330 milligrams per liter depending upon the degree of anesthetization or sedation desired, the species and size of the fish, and the temperature and softness of the water. Preliminary tests of solutions must be made with small numbers of fish to determine the desired rates of sedation or anesthesia and the appropriate exposure times for the specific lots of fish under prevailing conditions.